

ABLER_x



Affordable because we focus ONLY on equine medication

Ablver™

COMPOSITION

Each sachet contains:
Ivermectin BP..... 60 mg
(As film coated granules)

DESCRIPTION

Ablver™ granules of Ivermectin is a broad spectrum dewormer of easy-to-measure, easy-to-feed, flavourless, aqua coloured, film coated granules.

Ablver
product information

INDICATIONS

Ivermectin controls infestations due to adult and immature roundworms, lungworms and bots.

- Ascarids (adults and third- and fourth-stage larvae)
- Bots (oral and gastric larval stages)
- Hairworms (adults)
- Intestinal Threadworms (adults)
- Large Strongyles (adults) *Strongylus vulgaris*
- Large-mouth Stomach Worms (adults)
- Lungworms (adults and fourth-stage larvae)
- Pinworms (adults and fourth-stage larvae)
- Small Strongyles (adults, and some resistant to class of benzimidazoles)
- Small Strongyles (fourth-stage larvae)
- Summer Sores and Dermatitis caused by Neck Threadworm

DOSAGE

Control Program:

Recommended dosage: 0.2mg Ivermectin
200mg per 100kg (220lb) bodyweight. One sachet per 300kg (660lb) bodyweight
Consult your veterinarian for a control program to meet your specific needs. Regular treatment will reduce the chances of colic caused by verminous arteritis caused by *Strongylus vulgaris*.

ADMINISTRATION

Simply sprinkle Ablver™ dry onto feed. For fussy eaters mix with molasses or corn syrup then mix with feed, apple or a treat.

OVERDOSAGE

Should a massive overdose occur, treat symptomatically and supportively.

SIDE EFFECTS

Horses carrying heavy infections of onchocera microfilaries may have reactions such as swelling and itching after treatment. It is assumed that these reactions are the result of destruction of large numbers of microflora. Consult your veterinarian if these signs persist.

INTERACTIONS OR CONTRAINDICATIONS

Do not use in foals under 4 weeks of age.

TOXICITY

Reports indicate the combination of Ivermectin and Silver Nightshade plant creates a severe toxic affect on horses.

WARNINGS

Ablver™ has been formulated specifically for use in horses. Other animals may be adversely affected by the concentration of Ivermectin, if they are allowed to ingest spillages of paste or have access to the used syringes. Do not use on horses intended for human consumption. Keep out of reach of children. Veterinary use only.

DISPOSAL

Dispose of unused product and containers in accordance with guidance from your local waste regulation authority. Dangerous to fish and aquatic life, do not contaminate water ways with the product or used containers.

STORAGE

Store in a cool dry atmosphere away from extreme heat.
Store/Transport at and 20° to 30°C.(68 to 86°F)

CONTACT:

Tel: Toll Free +1-855-419-8993 USA
Email: helpdesk@abler.com
Website www.abler.com

