



Affordable because we focus ONLY on equine medication

AblerQuant™ 6.42g

COMPOSITION

Each syringe contains;
Ivermectin BP..... 1.87%
Praziquantel BP... 14.03%

AblerQuant
product information

DESCRIPTION

AblerQuant™ is a broad spectrum single dose dewormer in easy-to-use oral syringe of green apple-flavored dosages.

INDICATIONS

Ivermectin controls infestations due to adult and immature roundworms, lungworms and bots.
Praziquantel controls infestations of tapeworms.

Ascarids (adults and third- and fourth-stage larvae)
Bots (oral and gastric larval stages)
Hairworms (adults)
Intestinal Threadworms (adults)
Large Strongyles (adults) <i>Strongylus vulgaris</i>
Large-mouth Stomach Worms (adults)
Lungworms (adults and fourth-stage larvae)
Pinworms (adults and fourth-stage larvae)
Small Strongyles (adults, and some resistant to class of benzimidazoles)
Small Strongyles (fourth-stage larvae)
Summer Sores and Dermatitis caused by Neck Threadworm
TapeWorm - <i>Anoplocephala perfoliata</i>

DOSAGE

Control Program:

Controls Tapeworms and infestations due to adult and immature roundworms and lungworms.

Recommended dosage is 1.07g paste per 100kg (220lb) of horses bodyweight.

0.2mg Ivermectin and 1.5mg Praziquantel per kg bodyweight

- One (1) syringe doses a horse up to 600kg (1320lb) bodyweight.

Consult your veterinarian for a control program to meet your specific needs. Regular treatment will reduce the chances of colic caused by *Anoplocephala perfoliata* and verminous arteritis caused by *Strongylus vulgaris*

ADMINISTRATION

Before administering AblerQuant™ make sure the horse's mouth is free of food.

1. Rotate the ring to the appropriate mark on the syringe for your horses weight.
2. Lock into place by rotating the plunger ring 1/4 turn.
3. Remove the nozzle cover.
4. Insert the nozzle of the syringe through the interdental space, and deposit the required amount of paste at the back of the tongue.
5. Immediately after administration, elevate the horse's head for a few seconds to ensure the dose is swallowed.

IMPORTANT: If any dose is lost or rejected, re-dosing is recommended. Note that shelf life after first immediate opening is 28 days. Replace nozzle cover after each use.

OVERDOSAGE

Massive overdosage is required to cause problems in other species but the toxic dosage in horses is uncertain.

SIDE EFFECTS

Horses carrying heavy infections of onchocera microfilaries may have reactions such as swelling and itching after treatment.

It is assumed that these reactions are the result of destruction of large numbers of microflora. Consult your veterinarian if these signs persist.

INTERACTIONS OR CONTRAINDICATIONS

AblerQuant™ may be used in foals 4 weeks of age and older. Pregnant or lactating mares, breeding stallions may be treated without adverse effects on their fertility.

TOXICITY

Reports indicate the combination of Ivermectin and Silver Nightshade plant creates a severe toxic affect on horses.

WARNINGS

AblerQuant™ has been formulated specifically for use in horses. Other animals may be adversely affected by the concentration of Ivermectin, if they are allowed to ingest spillages of paste or have access to the used syringes.

Do not use on horses intended for human consumption.

Keep out of reach of children.

Veterinary use only.

STORAGE

Store in a cool dry atmosphere away from extreme heat.

Store/Transport at 20° to 30°C.(68° to 86°F)

DISPOSAL

Dispose of unused product and containers in accordance with guidance from your local waste regulation authority. Dangerous to fish and aquatic life, do not contaminate water ways with the product or used containers.

CONTACT

Tel: Toll Free +1-855-419-8993 USA

Email: helpdesk@abler.com

Website www.abler.com

